

## University of Groningen

### Time and age trends in smoking cessation in Europe

Ageing Lungs European Cohorts ALEC; Pesce, Giancarlo; Marcon, Alessandro; Calciano, Lucia; Perret, Jennifer L.; Abramson, Michael J.; Bono, Roberto; Bousquet, Jean; Fois, Alessandro G.; Janson, Christer

*Published in:*  
 PLoS ONE

*DOI:*  
 [10.1371/journal.pone.0211976](https://doi.org/10.1371/journal.pone.0211976)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
 Publisher's PDF, also known as Version of record

*Publication date:*  
 2019

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Ageing Lungs European Cohorts ALEC, Pesce, G., Marcon, A., Calciano, L., Perret, J. L., Abramson, M. J., Bono, R., Bousquet, J., Fois, A. G., Janson, C., Jarvis, D., Jogi, R., Leynaert, B., Nowak, D., Schlunssen, V., Urrutia-Landa, I., Verlato, G., Villani, S., Zuberbier, T., ... Accordini, S. (2019). Time and age trends in smoking cessation in Europe. *PLoS ONE*, 14(2), [0211976]. <https://doi.org/10.1371/journal.pone.0211976>

#### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

#### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

RESEARCH ARTICLE

# Time and age trends in smoking cessation in Europe

Giancarlo Pesce<sup>1,2,3\*</sup>, Alessandro Marcon<sup>1\*</sup>, Lucia Calciano<sup>1</sup>, Jennifer L. Perret<sup>4</sup>, Michael J. Abramson<sup>5</sup>, Roberto Bono<sup>6</sup>, Jean Bousquet<sup>7</sup>, Alessandro G. Fois<sup>8</sup>, Christer Janson<sup>9</sup>, Deborah Jarvis<sup>10,11</sup>, Rain Jögi<sup>12</sup>, Bénédicte Leynaert<sup>2,3</sup>, Dennis Nowak<sup>13</sup>, Vivi Schlünssen<sup>14</sup>, Isabel Urrutia-Landa<sup>15</sup>, Giuseppe Verlato<sup>1</sup>, Simona Villani<sup>16</sup>, Torsten Zuberbier<sup>17</sup>, Cosetta Minelli<sup>10</sup>, Simone Accordini<sup>10</sup>, on behalf of The Ageing Lungs in European Cohorts (ALEC) study<sup>1</sup>



## OPEN ACCESS

**Citation:** Pesce G, Marcon A, Calciano L, Perret JL, Abramson MJ, Bono R, et al. (2019) Time and age trends in smoking cessation in Europe. PLoS ONE 14(2): e0211976. <https://doi.org/10.1371/journal.pone.0211976>

**Editor:** Lamberto Manzoli, Università degli Studi di Ferrara, ITALY

**Received:** December 5, 2018

**Accepted:** January 24, 2019

**Published:** February 7, 2019

**Copyright:** © 2019 Pesce et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the manuscript and its Supporting Information files.

**Funding:** This study was conducted within the Ageing Lungs in European Cohorts study ([www.alecstudy.org](http://www.alecstudy.org)) and has received funding from the European Union's Horizon 2020 research and innovation programme (<https://ec.europa.eu/programmes/horizon2020/>) under grant agreement no 633212 (DJ). National funders who supported data collection in the original studies are listed in S1 Appendix. The funders had no role in study

**1** Unit of Epidemiology and Medical Statistics, Department of Diagnostics and Public Health, University of Verona, Verona, Italy, **2** UMR1152, Department of Pathophysiology and Epidemiology of Respiratory Diseases, INSERM, Paris, France, **3** UMR1152, Faculté de Médecine, Site Bichat, Université Paris-VII (Paris Diderot), Sorbonne Paris-Cité, Paris, France, **4** Allergy and Lung Health Unit, School of Population and Global Health, University of Melbourne, Melbourne, Australia, **5** Department of Epidemiology and Preventive Medicine, Monash University, Melbourne, Australia, **6** Department of Public Health and Pediatrics, University of Turin, Turin, Italy, **7** University Hospital and INSERM, Hôpital Arnaud de Villeneuve, Montpellier, France, **8** Clinical and Interventional Pulmonology, Department of Clinical, Surgical and Experimental Sciences, University of Sassari, Sassari, Italy, **9** Department of Medical Sciences: Respiratory, Allergy and Sleep research, Uppsala University, Uppsala, Sweden, **10** National Heart and Lung Institute, Imperial College London, London, United Kingdom, **11** MRC-PHE Centre for Environment and Health, Imperial College London, London, United Kingdom, **12** Department of Pneumology, University of Tartu, Tartu, Estonia, **13** Institute and Outpatient Clinic for Occupational, Social and Environmental Medicine, University Hospital of Munich (LMU), Munich, Germany, **14** Department of Public Health, Environment, Occupation and Health, Aarhus University, Aarhus, Denmark, **15** Respiratory Department, Galdakao Hospital, OSI Barrualde-Galdakao, Biscay, Spain, **16** Unit of Biostatistics and Clinical Epidemiology, Department of Public Health, Experimental and Forensic Medicine, University of Pavia, Pavia, Italy, **17** Allergy Centre Charité, Department of Dermatology & Allergy, Charité, Universitätsmedizin Berlin, Berlin, Germany

☞ These authors contributed equally to this work.

¶ Principal Investigators and Partners of the ALEC study are provided in the Acknowledgments

\* [giancarlopesce1981@gmail.com](mailto:giancarlopesce1981@gmail.com) (GP); [alessandro.marcon@univr.it](mailto:alessandro.marcon@univr.it) (AM)

## Abstract

### Background

Smoking is the main risk factor for most of the leading causes of death. Cessation is the single most important step that smokers can take to improve their health. With the aim of informing policy makers about decisions on future tobacco control strategies, we estimated time and age trends in smoking cessation in Europe between 1980 and 2010.

### Methods

Data on the smoking history of 50,228 lifetime smokers from 17 European countries were obtained from six large population-based studies included in the Ageing Lungs in European Cohorts (ALEC) consortium. Smoking cessation rates were assessed retrospectively, and age trends were estimated for three decades (1980–1989, 1990–1999, 2000–2010). The analyses were stratified by sex and region (North, East, South, West Europe).

design, data collection and analysis, decision to publish, or preparation of the manuscript. The corresponding author had full access to all the data and had final responsibility for the decision to submit for publication.

**Competing interests:** DJ reports research grants from the European Union and Medical Research Council. JLP reports a travel grant from the Thoracic Society of Australia and New Zealand (TSANZ) and Boehringer Ingelheim. MA reports research grants and/or personal fees from Pfizer, Boehringer-Ingelheim, Sanofi. TZ has received industry consulting, research grants and/or honoraria from: AstraZeneca, AbbVie, Almirall, Astellas, Bayer Health Care, Bencard, Berlin Chemie, FAES, HAL, Leti, Meda, Menarini, Merck, MSD, Novartis, Pfizer, Sanofi, Stallergenes, Takeda, Teva, UCB, Henkel, Kryolan, L'Oréal. TZ reports the following organizational affiliations: Committee member, WHO-Initiative "Allergic Rhinitis and its Impact on Asthma" (ARIA); Member of the Board, German Society for Allergy and Clinical Immunology (DGAKI); Head, European Centre for Allergy Research Foundation (ECARF); Secretary General, Global Allergy and Asthma European Network (GA2LEN); Member, Committee on Allergy Diagnosis and Molecular Allergology, World Allergy Organisation (WAO). VS reports research grants from The Wood Dust Foundation (project no. 444508795). All the other authors report no financial relationships with any organisations that might have an interest in the submitted work in the previous three years, and no other relationships or activities that could appear to have influenced the submitted work. These competing interests do not alter the authors' adherence to PLOS ONE policies on sharing data and materials. There are no restrictions to the sharing of data from this study.

**Abbreviations:** ALEC, Ageing Lungs In European Cohorts; ECRHS, European Community Respiratory Health Survey; GA2LEN, Global Allergy and Asthma Network of Excellence; GEIRD, Gene Environment Interactions in Respiratory Diseases; IRR, Incidence Rate Ratio; ISAYA, Italian Study on Asthma in Young Adults; RHINE, Respiratory Health in Northern Europe.

## Results

Overall, 21,735 subjects (43.3%) quit smoking over a total time-at-risk of 803,031 years. Cessation rates increased between 1980 and 2010 in young adults (16–40 years), especially females, from all the regions, and in older adults (41–60 years) from North Europe, while they were stable in older adults from East, South and West Europe. In the 2000s, the cessation rates for men and women combined were highest in North Europe (49.9 per 1,000/year) compared to the other regions (range: 26.5–32.7 per 1,000/year). A sharp peak in rates was observed for women around the age of 30, possibly as a consequence of pregnancy-related smoking cessation. In most regions, subjects who started smoking before the age of 16 were less likely to quit than those who started later.

## Conclusions

Our findings suggest an increasing awareness on the detrimental effects of smoking across Europe. However, East, South and West European countries are lagging behind North Europe, suggesting the need to intensify tobacco control strategies in these regions. Additional efforts should be made to keep young adolescents away from taking up smoking, as early initiation could make quitting more challenging during later life.

## Introduction

Smoking is the main risk factor for most of the leading causes of death.[1] Every year about 1.6 million Europeans die because of smoking, making Europe the region with the greatest proportion of deaths (16%) attributable to tobacco worldwide.[2] Tobacco smoking is also a financial burden on European healthcare systems and economies.[3]

In developed countries, smoking has been associated with a loss of at least a decade of life.[4] However cessation, especially before the age of 40 years, can dramatically reduce the risk of death and significantly improve quality of life.[4] It is therefore critical to encourage all current smokers to quit in order to reduce the burden of smoking-related diseases and mortality.[2,3]

European countries have introduced various tobacco control policies (e.g. increased taxation of tobacco products, no-smoking areas, advertising bans, graphic pack warnings, and smoking cessation programs) over recent decades. Despite a general decrease in tobacco consumption and the prevalence of smoking, still more than one in four of the European population over the age of 14 years is an active smoker.[5]

Variations in smoking prevalence are determined by variations in the rates of initiation, cessation and relapse. For this reason, monitoring trends in these rates is fundamental for evaluating current tobacco control policies and for planning future strategies that can effectively tackle the tobacco epidemic.[2] Nonetheless, there is an extreme paucity of studies on smoking cessation in the general European population.

With the aim of better informing public health policy makers about decisions on future tobacco control strategies, we estimated age trends in smoking cessation in Europe for the three decades between 1980 and 2010. The present analysis compliments our previous report on smoking initiation in Europe [6].

## Methods

### Study design and population

We analysed data obtained from six multicentre studies on random samples of the general adult population carried out in Europe between 1991 and 2013, which became part of the Ageing Lungs in European Cohorts (ALEC) consortium ([www.alecstudy.org](http://www.alecstudy.org)).

The Global Allergy and Asthma Network of Excellence (GA<sup>2</sup>LEN, [www.ga2len.net](http://www.ga2len.net)) is an international cross-sectional study that was carried out on people aged 15–75 years between 2007 and 2009.[7] The Italian Study on Asthma in Young Adults (ISAYA) is a postal cohort study on subjects aged 20–44 at enrolment in 1998–2000, who were followed up in 2008–2009.[8] The Genes Environmental Interaction in Respiratory Diseases (GEIRD, [www.geird.org](http://www.geird.org)) is a cross-sectional study carried out in Italy between 2005 and 2010.[9]

The European Community Respiratory Health Survey (ECRHS, [www.ecrhs.org](http://www.ecrhs.org)) is an international cohort study performed on subjects aged 20–44 years at enrolment in 1991–1994.[10] In ECRHS I, random samples of subjects participating in a postal screening (stage 1) took part in a clinical interview (stage 2), where they reported information on smoking habits for the first time. The cohort of participants in stage 2 (labelled “ECRHS clinical” throughout the manuscript) was reassessed at two independent examinations during 1999–2002 in ECRHS II, [11] and during 2010–2013 in ECRHS III.[12] The Italian arm of the ECRHS (ECRHS-Italy) is the postal follow-up of the Italian participants in ECRHS stage-1, who were surveyed in 1998–2001 and 2008–2009.[13,14] The Respiratory Health in Northern Europe study (RHINE, [www.rhine.nu](http://www.rhine.nu)) is a postal follow-up of the ECRHS I stage-1 participants from Scandinavian and Estonian centres, carried out in 2010–2012 (RHINE II).[15] A flow-chart describing how the three cohorts stem from the ECRHS study can be found at <https://doi.org/10.1371/journal.pone.0201881.s009>.

The analyses were performed on data pooled from the six studies. Subjects who had participated in more than one study or occasion were only considered once. The distribution of subjects by sample (i.e. the original centre and study where the subjects were surveyed) is shown in [S1 Table](#). Overall, 73 samples from 53 centres in 17 countries were included, while four samples with a low participation rate (below 25%) were discarded. The samples were grouped into four regions, according to the United Nations geoscheme and tobacco epidemiology:[16] South Europe (Italy, Portugal, Spain), West Europe (Belgium, France, Germany, the Netherlands, Switzerland), North Europe (Denmark, Finland, Iceland, Norway, Sweden, United Kingdom), and East Europe (Estonia, Macedonia, Poland).

### Data on smoking

The smoking history of each subject was reconstructed retrospectively using data from postal questionnaires for all studies, in addition to data from clinical interviews for ECRHS.[[Table 1](#)] In all studies, with the exception of RHINE (where the subjects qualified themselves as “smokers” or “ex-smokers”), smoking was defined as active smoking for as long as a year (regular smoking). Life-long never smokers were excluded from the analyses. A smoker who reported not to have smoked at all in the last month was considered an ex-smoker. Age at initiation was based on the question “How old were you when you started smoking?” for all the studies. Questions on age at cessation referred to the time of stopping or the last time smoking, with a slightly different wording across studies.[[Table 1](#)] For subjects with follow-up data (ECRHS clinical, ECRHS-Italy, ISAYA), the first information available was chosen for defining age at initiation, while age at cessation was based on the last information available.

Table 1. Questionnaire items on smoking.

Study	Smoking status	Age at initiation	Current smoking	Age at cessation
GA <sup>2</sup> LEN	Have you ever smoked for as long as a year? ['YES' means at least one cigarette per day or one cigar per week for one year]	How old were you when you started smoking?	Have you smoked at all in the last month?	How old were you when you stopped smoking?
RHINE	Are you a smoker? (this applies even if you only smoke the odd cigarette/ cigar or pipe every week)		Are you an ex-smoker?	Stopped smoking in . . . (year)
ISAYA, ECRHS-Italy, GEIRD	Have you ever smoked for as long as a year? ['YES' means at least one cigarette per day or one cigar per week for one year]		Do you now smoke, as of one month ago?	How old were you when you stopped smoking?
ECRHS (clinical interview)	Have you ever smoked for as long as a year? ['YES' means at least 20 packs of cigarettes or 12 oz (360 grams) of tobacco in a lifetime, or at least one cigarette per day or one cigar per week for one year]		Do you now smoke, as of one month ago?	How old were you when you stopped or cut down smoking?*
ECRHS (postal questionnaire)	Have you ever smoked for as long as a year?		Have you smoked in the last month?	How old were you when you last smoked?

\*only the answers from subjects who quit smoking were considered.

<https://doi.org/10.1371/journal.pone.0211976.t001>

## Statistical analysis

Rates of smoking cessation (per 1,000/year) were calculated as the ratio between the number of incident quitters and the total time at risk, which was defined as years from initiation to cessation for quitters, or from initiation to the last survey available for active smokers. To avoid data sparseness, time at risk was left-censored at age 16 years and at 1980, and right-censored at age 60 years and at 2010.

All the analyses were stratified by sex, region, and period (1980–1989, 1990–1999, 2000–2010). Crude rates were calculated for two age groups: 16–40 and 41–60 years. In the main analyses, age-specific trends of smoking cessation rates (with 95% confidence intervals) were estimated using two-level mixed-effects models (with subjects nested into samples), with a negative binomial outcome distribution, a logarithmic link function and an offset for log(person-years). Period was included as a linear term to adjust for possible time trends of cessation rates within each decade. Age was modelled using natural smoothed polynomial lines (splines), and the number of knots (four and five for males and females, respectively) was chosen based on model fit using the Bayesian information criterion.

As the risk of relapse is high in the first year after quitting, while it drastically drops afterwards,[17] we performed a sensitivity analysis defining smoking cessation as having quit smoking for at least two years (with person-years re-calculated accordingly), in order to exclude subjects who might have quit smoking just for a brief time before the survey.

Finally, to evaluate whether subjects who started smoking in early adolescence had a different attitude toward smoking cessation compared to those who started later, the main models were fitted including a binary indicator for age at initiation (10–15 vs.  $\geq 16$  years) as a predictor variable. In this analysis, since we found no evidence of interaction between age at initiation and period, the three decades were combined to increase the statistical power.

The statistical analyses were performed using STATA 14.2 (StataCorp, College Station, TX).

## Ethics approval and consent to participate

Ethical approval was not requested for this secondary analysis of pooled data from previous studies. In each of the original studies, ethical approval was obtained for each centre from the appropriate ethics committee. All procedures have conformed to the principles embodied in

the Declaration of Helsinki. Written informed consent was obtained from participants in the clinical examinations (ECRHS clinical). In ECRHS-Italy, RHINE, GA<sup>2</sup>LEN, ISAYA and GEIRD, only data from postal questionnaires were used, which were voluntarily sent back giving consent to use the anonymized data.

## Results

A total of 50,228 ever smokers with complete information on sex, year of birth, age at smoking initiation and, if applicable, age at cessation were included in the analysis.[S1 Fig and S1 Table] The distribution of the participants by study and year is shown in S2 Fig. GA<sup>2</sup>LEN was the study that contributed the highest number of subjects (n = 22,100). Overall, 14.0% (n = 7,058) of the subjects had follow-up data.

Females represented about half of the sample, and the median age at the last interview ranged from 38 to 50 years for subjects from South and North Europe, respectively. Overall, 43.3% (n = 21,735) of the subjects quit smoking over a total time-at-risk of 803,031 person-years. The proportion of quitters was highest in North Europe (56.0%) compared to other European regions (range: 34.1% to 37.9%).[Table 2]

## Trends in smoking cessation

In the 2000s, the crude cessation rates for males and females combined were highest in North Europe (49.9 per 1,000/year) compared to the other regions (range: 26.5–32.7 per 1,000/year). For males and females aged less than 40 years, cessation rates were considerably higher in the 2000s compared to the preceding two decades in all the four regions. This was especially evident in North Europe, where the cessation rates doubled, from 24 per 1,000/year in the 1980s to 53 per 1,000/year in the 2000s. On the other hand, cessation rates after the age of 40 were similar across successive periods, with the exception of North Europe where the rates doubled, from 23 per 1,000/year in the 1980s to 47 per 1,000/year in the 2000s.

Consistent results were found when looking separately at males and females.[Table 3]

Table 2. Demographic and smoking history data of the subjects included in the analyses, by region.

	North Europe	East Europe	South Europe	West Europe	Overall
Countries, n	6	3	3	5	17
Centres, n	15	5	19	14	53
Subjects included, n	18,255	5,013	19,179	7,781	50,228
Females, n (%)	9,927 (54.4)	2,399 (47.9)	8,716 (45.5)	3,882 (49.9)	24,924 (49.6)
Birth year *	1958 (1940–1983)	1961 (1940–1986)	1964 (1948–1980)	1961 (1943–1986)	1962 (1942–1984)
Year of last interview *	2008 (2002–2011)	2008 (2007–2012)	2001 (1998–2010)	2008 (1991–2012)	2008 (1998–2011)
Age at last interview (years) *	50 (24–68)	47 (22–67)	38 (23–60)	43 (22–66)	43 (23–66)
Study, n (%)					
ECRHS clinical	2,389 (13.1)	214 (4.3)	1,764 (9.2)	3,263 (41.9)	7,630 (15.2)
ECRHS-Italy	-	-	1,808 (9.4)	-	1,808 (3.6)
RHINE	3,665 (20.1)	364 (7.3)	-	-	4,029 (8.0)
ISAYA	-	-	8,788 (45.8)	-	8,788 (17.5)
GEIRD	-	-	5,873 (30.6)	-	5,873 (11.7)
GA <sup>2</sup> LEN	12,201 (66.8)	4,435 (88.5)	946 (4.9)	4,518 (58.1)	22,100 (44.0)
Quitters, n (%)	10,224 (56.0)	1,707 (34.1)	6,854 (35.7)	2,950 (37.9)	21,735 (43.3)
Total person-years	309,077	89,194	287,637	117,123	803,031

\* median (5<sup>th</sup>–95<sup>th</sup> percentile)

<https://doi.org/10.1371/journal.pone.0211976.t002>



Table 3. Crude rates of smoking cessation (per 1,000/year), number of quitters and person-years at risk in males and females by region, age group and period.

Region	Age group		Males			Females		
			1980–1989	1990–1999	2000–2010	1980–1989	1990–1999	2000–2010
North Europe	16–40 years	rates†	24.7	31.2	49.8	23.0	33.4	56.1
		quitters	1,150	950	714	1,252	1,321	1,111
		py	46,594	30,445	14,336	54,329	39,553	19,808
	41–60 years	rates†	25.0	31.1	48.6	20.3	28.7	45.6
		quitters	227	667	979	171	623	1,059
		py	9,066	21,422	20,131	8,444	21,733	23,216
East Europe	16–40 years	rates†	11.8	16.8	23.5	9.4	15.1	30.2
		quitters	151	190	172	105	153	200
		py	12,845	11,312	7,320	11,210	10,146	6,619
	41–60 years	rates†	22.4	28.9	28.5	18.0	19.5	24.3
		cases	60	175	190	31	108	172
		py	2,682	6,051	6,665	1,722	5,541	7,081
South Europe	16–40 years	rates†	17.8	22.4	33.3	20.4	24.6	36.9
		quitters	961	1,376	649	930	1,256	610
		py	54,084	61,528	19,513	45,515	51,002	16,547
	41–60 years	rates†	30.3	30.1	33.8	15.1	21.2	22.2
		quitters	94	299	317	20	149	193
		py	3,098	9,946	9,373	1,323	7,015	8,693
West Europe	16–40 years	rates†	20.8	24.1	33.3	22.3	22.8	32.7
		quitters	459	304	253	461	295	300
		py	22,118	12,608	7,609	20,678	12,951	9,162
	41–60 years	rates†	32.6	28.0	32.4	22.9	24.9	23.1
		cases	93	193	240	43	140	169
		py	2,856	6,891	7,416	1,878	5,633	7,323
Overall	16–40 years	rates†	20.1	24.3	36.7	20.9	26.6	42.6
		quitters	2,721	2,820	1,788	2,748	3,025	2,221
		py	135,641	115,893	48,778	131,732	113,652	52,136
	41–60 years	rates†	26.8	30.1	39.6	19.8	25.5	34.4
		cases	474	1,334	1,726	265	1,020	1,593
		py	17,702	44,310	43,585	13,367	39,922	46,313


†: quitters per 1,000/year. py: person-years.

<https://doi.org/10.1371/journal.pone.0211976.t003>

Fig 1 shows the smoothed age-trends in smoking cessation by sex, region, and period.[Fig 1] We observed a peak in smoking cessation around the age of 30, which was particularly evident in the 2000s. This peak was quite blunted in men, while it was very sharp in women and corresponded to an age of 31 years in South Europe, 28 years in West and North Europe, and 26 years in East Europe. From the age of 40 onwards, the smoking cessation rates increased gradually in males and females from all regions.

### Smoking cessation according to age at initiation


Except for males from South Europe, where the incidence rate ratio (IRR) for later age at initiation was below 1 and not statistically significant (IRR 0.97, 95%CI: 0.90–1.04), the likelihood of quitting was higher for both men and women who started smoking at 16 years or later compared to those who started earlier, with IRRs ranging from 1.05 (95%CI: 0.88–1.24) in East European males to 1.14 (95%CI: 1.07–1.21) in North European males.[Fig 2]



**Fig 1. Estimated trends in smoking cessation rates (per 1,000/year) with 95% confidence intervals in females and males, by region.** Trends were stratified by period: black lines refer to 1980–1989; red lines refer to 1990–1999, and green lines refer to 2000–2010.

<https://doi.org/10.1371/journal.pone.0211976.g001>





**Fig 2.** Age and period-adjusted incidence rate ratios of smoking cessation, with 95% confidence intervals, for subjects who started smoking at 16 years or later compared to those who started earlier, by sex and European region.

<https://doi.org/10.1371/journal.pone.0211976.g002>

### Probability of smoking relapse after quitting

Among ex-smokers, the median duration of abstinence from smoking reported at the time of the interview was 10 years (25<sup>th</sup>-75<sup>th</sup> percentiles: 4–18), and over 90% had quit for at least 2 years.

The risk of relapse was estimated using subjects with follow-up data.[S2 Table] Overall, 12.2% of those who were ex-smokers at the baseline examination reported to be active smokers at the last follow-up (average follow-up time: 16.5 years). The proportion of subjects relapsing was significantly greater in those who had quit smoking for less than 2 years at baseline (nearly 30%) compared to those who had quit smoking for at least 2 years (less than 10%) ( $p < 0.001$ ).

### Discussion

We retrospectively investigated trends in smoking cessation across three decades from 1980 to 2010, using representative samples of the general population from 17 European countries.

The main results of our analyses are the following:

- rates of smoking cessation were higher in North Europe than in West, South, and East Europe;

- smoking cessation has been steadily increasing between the 1980s and the 2000s among smokers aged under 40 years across Europe; among older people, this was observed only for North Europe;
- smoking cessation rates peak around the age of 30 in women;
- quitting was less likely for subjects who took up smoking before the age of 16 compared to people starting later.

### Time trends in smoking cessation across European regions

We showed that the rate of smoking cessation was 49.9 per 1,000/year in North Europe (United Kingdom and Scandinavian countries) in 2000–2010, which is about 60% higher compared to the other regions. Cessation rates for North Europe confirm the rates recently reported from RHINE, which is one of the cohorts included in this study,[18] while we are not aware of studies reporting cessation rates for the other regions.

We found that the rates of cessation have been increasing in North Europe since the end of the last century, while in the rest of Europe they have started to grow only after the year 2000. Moreover, in North Europe, the increasing trends in smoking cessation involved both young (16–40 years) and older (41–60 years) adults, while in the rest of Europe they were restricted to young adults. These results compliment a previous report on smoking initiation using data from the same studies, where we observed lower rates of uptake of regular smoking in North Europe compared to the other regions.[6] Overall, the geographical pattern in smoking cessation described in our study is consistent with the pattern in smoking prevalence across European countries reported by others [4, 19,20]. These data testify to the better implementation of effective tobacco control policies in countries of North Europe.[21] Of note, the higher attained education and wealth per-capita in North Europe might also have contributed to boost tobacco control strategies, as preventative measures are more effective among higher socioeconomic groups.[22]

### Age trends in smoking cessation in males and females

We found that the rates of smoking cessation were broadly similar in men and women within the same European region in the 2000s, but there was a remarkable gender difference in age trends. In fact, while the cessation rates were similar in young and older men, women aged less than 40 years had significantly higher rates than older women. These findings are in agreement with another study on population data from USA, Canada, and UK, which showed that women were more likely to have given up smoking than men before their fifties, and *vice-versa* after the age of 50.[23]

We also found a sharp peak of cessation around the age of 30 in women from all regions, which was especially evident in the 2000s. The peak slightly shifted at different ages according to the region, matching well with mothers' mean age at first child that ranges from 26–27 years in East Europe to 30–31 years in South Europe.[24] This suggests that pregnancy is the primary reason for quitting in women. Previous reports showed that about three out of 4 women stop smoking during pregnancy,[25] and cessation is more frequent when expecting the first child.[26]

The more evident peaks of cessation in the 2000s most likely reflect an increasing awareness of the negative health impact associated with smoking during pregnancy for both the mother and the foetus,[27] and growing perception of the need to protect children from second-hand smoke exposure. Social stigma towards expecting mothers who smoke can also encourage

cessation.[28] The blunted peak observed around the age of 30 among males may also be determined by parenthood, as fathers try to abstain from smoking to be supportive to their partners' attempts to quit during pregnancy,[29,30] and to avoid their children being exposed to second-hand smoke.[31]

After the age of 40, rates of cessation gradually increased in both men and women, maybe indicating a growing concern about smoking-related illnesses with ageing.[32]

### Age at initiation affects the likelihood to quit

Although overall rates of smoking initiation have decreased considerably in Europe during recent decades, initiation in young adolescents (11–15 years) has alarmingly increased.[6] Early initiation is associated with increased risk of smoking-related mortality and morbidity.[33] Males starting smoking before age 15 may have children with an increased risk of respiratory diseases, possibly through epigenetic changes in sperm precursor cells.[34] We found that early age at initiation is associated with lower rates of cessation, in agreement with other studies from North America.[35,36] Early initiation has been associated with greater risk of relapse after cessation.[37] Younger adolescents develop nicotine addiction after a shorter duration of smoking and a lower number of cigarettes compared to those who take up smoking later.[38] All these findings suggest that tobacco control strategies targeting the youngest should be implemented with the aim of discouraging early smoking, or at least delaying initiation.

### Strengths and limitations

The main strengths of our study are the inclusion of large samples of subjects from 17 European countries and a long observation period, allowing the comparison of cessation rates across different regions and their trends during three decades.

A drawback is that this is an observational retrospective study that relies on self-reported information on smoking history, and smoking status was not verified with biochemical methods. However, the agreement between self-reported smoking and cotinine levels is generally good in population studies.[39]

Another limitation is the potential telescoping bias, as individuals might be inaccurate in recalling the timing of past events. However, we found that age at smoking initiation was consistently reported up to two decades apart (see <https://doi.org/10.1371/journal.pone.0201881.s001> and S3 Table).

The risk of relapse is the main issue when studying smoking cessation. About half of smokers attempts to quit every year, but less than 10% achieves abstinence for a period of at least 6 months.[40] The first year after a quit attempt is the period at highest risk for relapse.[17] In our study, age at cessation reported in a sub-sample of ex-smokers with follow-up data (ECRHS clinical) increased by two years on average after 20 years, probably indicating smoking relapses. However, we derived age at cessation from the last time point available, and over 90% of quitters had abstained from smoking for at least two years before the interview, suggesting that our findings mainly reflect smoking cessation over an extended period among subjects at a low risk of relapse. Moreover, the main results were confirmed in a sensitivity analysis using a more conservative definition of smoking cessation (having quit for at least two years; S3 Fig).

### Conclusions

Our findings indicate that smoking cessation rates are increasing throughout Europe, especially for women around the age of 30, suggesting that there is a mounting awareness of the detrimental effects of smoking exposure. Scandinavian countries and the UK have greater

cessation rates than the rest of Europe, where the implementation of tobacco control strategies should be intensified. Additional efforts should be made to prevent adolescents taking up smoking, as early initiation could make quitting smoking more challenging during later life.

## Supporting information

**S1 Appendix. Supplementary information on the original studies.**  
(DOCX)

**S1 Dataset. Minimal dataset to replicate the analyses.** The dataset includes: region (North, East, South, West Europe); sex (male = 1, female = 2), a = age in years, p = period (year of observation), c = cohort (birth year), d = number of quitters, py = person-years.  
(CSV)

**S1 Fig. Study participants.**  
(DOCX)

**S2 Fig. Distribution of ever smokers by study and year of interview.**  
(DOCX)

**S3 Fig. Sensitivity analysis. Estimated trends in the rates (per 1,000/year) of smoking cessation (defined as having quitted smoking for at least two years) with 95% confidence intervals in females and males, by region.** Trends were stratified by period: black lines refer to 1980–1989; red lines refer to 1990–1999, and green lines refer to 2000–2010.  
(DOCX)

**S1 Table. Distribution of ever smokers included in the analysis (and total number of participants into the original studies with complete data) by region, centre and study.**  
(DOCX)

**S2 Table. Cumulative incidence of smoking relapse by time since quitting.**  
(DOCX)

**S3 Table. Comparison of the age at smoking initiation and cessation reported at different occasions in multi-wave studies.**  
(DOCX)

## Acknowledgments

The ALEC study leader is Deborah Jarvis. The manuscript was done under ALEC Workpackage 6 led by Cosetta Minelli. Other Workpackage leaders in ALEC are Cecilie Svanes, John Henderson (Department of Community Based Medicine, University of Bristol, Bristol, UK), Judith Garcia-Aymerich, Nicole Probst-Hensch (Swiss Tropical and Public Health Institute and University of Basel, Basel, Switzerland). The principal investigators and team members of the original studies are reported in [S1 Appendix](#) and in the study website <https://www.alecstudy.org/>.

The ALEC International Scientific Advisory Board is: Marike Boezen (University Medical Center Groningen, University of Groningen, Groningen, The Netherlands); Bernice Elger (Institute for Biomedical Ethics, University of Basel, Basel, Switzerland); Bo Alexander Gleditsch (The Norwegian Asthma and Allergy Association, Norway); Bas Heijmans (Department of Medical Statistics and Bioinformatics, Leiden University Medical Center, Leiden, The Netherlands); Isabelle Romieu (National Institute of Public Health, Cuernavaca, Mexico; and

Emory University, Atlanta, US); John Thompson (Department of Health Sciences, University of Leicester, Leicester, UK).

## Author Contributions

**Conceptualization:** Giancarlo Pesce, Alessandro Marcon, Cosetta Minelli, Simone Accordini.

**Data curation:** Giancarlo Pesce, Alessandro Marcon.

**Formal analysis:** Giancarlo Pesce, Alessandro Marcon, Lucia Calciano, Simone Accordini.

**Funding acquisition:** Deborah Jarvis.

**Investigation:** Giancarlo Pesce, Alessandro Marcon, Simone Accordini.

**Methodology:** Giancarlo Pesce, Alessandro Marcon, Lucia Calciano, Cosetta Minelli, Simone Accordini.

**Resources:** Michael J. Abramson, Roberto Bono, Jean Bousquet, Alessandro G. Fois, Christer Janson, Deborah Jarvis, Rain Jögi, Bénédicte Leynaert, Dennis Nowak, Vivi Schlünssen, Isabel Urrutia-Landa, Giuseppe Verlato, Simona Villani, Torsten Zuberbier, Simone Accordini.

**Supervision:** Alessandro Marcon, Simone Accordini.

**Writing – original draft:** Giancarlo Pesce.

**Writing – review & editing:** Giancarlo Pesce, Alessandro Marcon, Lucia Calciano, Jennifer L. Perret, Michael J. Abramson, Roberto Bono, Jean Bousquet, Alessandro G. Fois, Christer Janson, Deborah Jarvis, Rain Jögi, Bénédicte Leynaert, Dennis Nowak, Vivi Schlünssen, Isabel Urrutia-Landa, Giuseppe Verlato, Simona Villani, Torsten Zuberbier, Cosetta Minelli, Simone Accordini.

## References

1. Carter MPH, Abnet CC, Feskanich D, Freedman ND, Hartge P, Lewis CE, et al. Smoking and Mortality—Beyond Established Causes. *N Eng J Med* 2015; 372:631–640.
2. World Health Organization (WHO) 2017. Report on the global tobacco epidemic, 2017. Monitoring tobacco use and prevention policies. Geneva: World Health Organization 2017. (<http://apps.who.int/iris/bitstream/10665/255874/1/9789241512824-eng.pdf?ua=1&ua=1>, accessed 1 October 2018)
3. European Respiratory Society (ERS). The European Lung White Book. Chapter 9: Major risk factors—Tobacco Smoking. ([https://www.erswhitebook.org/files/public/Chapters/09\\_tobacco\\_smoking.pdf](https://www.erswhitebook.org/files/public/Chapters/09_tobacco_smoking.pdf), accessed 1 October 2018).
4. Jha P, Ramasundarahettige C, Landsman V, Rostron B, Thun M, Anderson RN, et al. 21st-century hazards of smoking and benefits of cessation in the United States. *N Engl J Med* 2013; 368:341–350. <https://doi.org/10.1056/NEJMs1211128> PMID: 23343063
5. Eurobarometer. Special Report 458: Attitudes of Europeans toward tobacco and electronic cigarettes. May 2017. (<https://ec.europa.eu/commfrontoffice/publicopinion/index.cfm/ResultDoc/download/DocumentKy/79002>, accessed 1 October 2018)
6. Marcon A, Pesce G, Calciano L, Bellisario V, Dharmage SC, Garcia-Aymerich J, et al. Trends in smoking initiation in Europe over 40 years: a retrospective cohort study. *PLoS One* 2018; 13(8):e0201881 <https://doi.org/10.1371/journal.pone.0201881> PMID: 30133533
7. Jarvis D, Newson R, Lotvall J, Hastan D, Tomassen P, Keil T, et al. Asthma in adults and its association with chronic rhinosinusitis: the GA2LEN survey in Europe. *Allergy* 2012; 67:91–8. <https://doi.org/10.1111/j.1398-9995.2011.02709.x> PMID: 22050239
8. de Marco R, Poli A, Ferrari M, Accordini S, Giammanco G, Bugiani M, et al. The impact of climate and traffic-related NO<sub>2</sub> on the prevalence of asthma and allergic rhinitis in Italy. *Clin Exp Allergy* 2002; 32:1405–1412. PMID: 12372117
9. de Marco R, Accordini S, Antonicelli L, Bellia V, Bettin MD, Bombieri C, et al. The Gene-Environment Interactions in Respiratory Diseases (GEIRD) project. *Int Arch Allergy Immunol* 2010; 153(3):255–63.

10. Burney PG, Luczynska C, Chinn S, Jarvis D. The European Community Respiratory Health Survey. *Eur Respir J* 1994; 7:954–60. PMID: [8050554](#)
11. European Community Respiratory Health Survey II Steering Committee. The European Community Respiratory Health Survey II. *Eur Respir J* 2002; 20:1071–9. PMID: [12449157](#)
12. Amaral AFS, Newson RB, Abramson MJ, Antó JM, Bono R, Corsico AG, et al. Changes in IgE sensitization and total IgE levels over a 20 years follow-up. *J Allergy Clin Immunol* 2016; 137(6):1788–95. <https://doi.org/10.1016/j.jaci.2015.09.037> PMID: [26586040](#)
13. de Marco R, Cappa V, Accordini S, Rava M, Antonicelli L, Bortolami O, et al. Trends in the prevalence of asthma and allergic rhinitis in Italy between 1991 and 2010. *Eur Respir J* 2012; 39:883–92. <https://doi.org/10.1183/09031936.00061611> PMID: [22005911](#)
14. Pesce G, Locatelli F, Cerveri I, Bugiani M, Pirina P, Johannessen A, et al. Seventy years of asthma in Italy: age, period and cohort effects on incidence and remission of self-reported asthma from 1940 to 2010. *PLoS One* 2015; 10(10):e0138570. <https://doi.org/10.1371/journal.pone.0138570> PMID: [26439263](#)
15. Toren K, Gislason T, Omenaas E, Jögi R, Forsberg B, Nyström L, et al. A prospective study of asthma incidence and its predictors: the RHINE study. *Eur Respir J* 2004; 24(6):942–6. <https://doi.org/10.1183/09031936.04.00044804> PMID: [15572536](#)
16. Bilano V, Gilmour S, Moffiet T, d'Espaignet ET, Stevens GA, Commar A, et al. Global trends and projections for tobacco use, 1990–2025: an analysis of smoking indicators from the WHO Comprehensive Information Systems for Tobacco Control. *Lancet* 2015; 385:966–76. [https://doi.org/10.1016/S0140-6736\(15\)60264-1](https://doi.org/10.1016/S0140-6736(15)60264-1) PMID: [25784347](#)
17. Garcia-Rodriguez O, Secades-Villa R, Florez-Salamanca L, Okuda M, Liu SM, Blanco C. Probability and predictors of relapse to smoking: results of the National Epidemiologic Survey on Alcohol and Related Conditions (NESARC). *Drug Alcohol Depend* 2013; 132:479–85. <https://doi.org/10.1016/j.drugalcdep.2013.03.008> PMID: [23570817](#)
18. Holm M, Schioler L, Andersson E, Forsberg B, Gislason T, Janson C, et al. Predictors of smoking cessation: a longitudinal study in a large cohort of smokers. *Respir Med* 2017; 123:164–9.
19. Bogdanovica I, Godfrey F, McNeill A, Britton J. Smoking prevalence in the European Union: a comparison of national and transnational prevalence survey methods and results. *Tob Control* 2011; 20:e4. <https://doi.org/10.1136/tc.2010.036103> PMID: [20966129](#)
20. Zatonski W, Przewoniak K, Sulkowska U, West R, Wojtyla A. Tobacco smoking in countries of the European Union. *Ann Agricul Environ Med* 2012; 19(2):181–192.
21. Joossens L, Raw M. The Tobacco Control Scale: a new scale to measure country activity. *Tobacco Control* 2006; 15:247–253. <https://doi.org/10.1136/tc.2005.015347> PMID: [16728757](#)
22. Bosdriesz JR, Willemsen MC, Stronks K, Kunst AE. Tobacco control policy and socio-economic inequalities in smoking in 27 European countries. *Drug Alcohol Dependence* 2016; 165:79–86 <https://doi.org/10.1016/j.drugalcdep.2016.05.020> PMID: [27262899](#)
23. Jarvis MJ, Cohen JE, Delnevo CD, Giovino GA. Dispelling myths about gender differences in smoking cessation: population data from USA, Canada, and Britain. *Tob Control* 2013; 22:356–60. <https://doi.org/10.1136/tobaccocontrol-2011-050279> PMID: [22649182](#)
24. Central Intelligence Agency (CIA). The World Factbook—Mother Mean Age at First Child. (<https://www.cia.gov/library/publications/the-world-factbook/fields/2256.html>, accessed 1 October 2018)
25. Smedberg J, Lupattelli A, Mardby AC, Nordeng H. Characteristics of women who continue smoking during pregnancy: a cross-sectional study of pregnant women and new mothers in 15 European countries. *BMC Pregnancy Childbirth* 2014; 14:213. <https://doi.org/10.1186/1471-2393-14-213> PMID: [24964728](#)
26. Tsakiridis I, Mamopoulos A, Papazisis G, Petousis S, Liozidou A, Athanasiadis A, et al. Prevalence of smoking during pregnancy and associated risk factors: a cross-sectional study in Northern Greece. *Eur J Pub Health* 2018; 28(2):321–5.
27. Einarson A, Riordan S. Smoking in pregnancy and lactation: a review of risks and cessation strategies. *Eur J Clin Pharmacol* 2009; 65:325–30. <https://doi.org/10.1007/s00228-008-0609-0> PMID: [19169678](#)
28. Wigginton B, Lee C. Stigma and hostility towards pregnant smokers: does individuating information reduce the effect? *Psychol Health* 2013; 28(8):862–73 <https://doi.org/10.1080/08870446.2012.762101> PMID: [23343130](#)
29. McBride CM, Baucom DH, Peterson BL, Pollak KI, Palmer C, Westman E, et al. Prenatal and postpartum smoking abstinence: a partner-assisted approach. *Am J Prev Med* 2004; 27(3):232–8. <https://doi.org/10.1016/j.amepre.2004.06.005> PMID: [15450636](#)
30. Janson C, Kunzli N, de Marco R, Chinn S, Jarvis D, Svanes C, et al. Changes in active and passive smoking in the European Community Respiratory Health Survey. *Eur Respir J* 2006; 27:517–24. <https://doi.org/10.1183/09031936.06.00106605> PMID: [16507851](#)



31. Flemming K, Graham H, McCaughan D, Angus K, Bauld L. The barriers and facilitators to smoking cessation experienced by women's partners during pregnancy and the post-partum: a systematic review of qualitative research. *BMC Public Health* 2015; 15:849. <https://doi.org/10.1186/s12889-015-2163-x> PMID: 26335935
32. Coombs RB, Li S, Kozlowski LT. Age interacts with heaviness of smoking in predicting success in cessation of smoking. *Am J Epidemiol* 1992; 135:240–6. PMID: 1546699
33. Choi SH, Stommel M. Impact of age at smoking initiation on smoking-related morbidity and all-cause mortality. *Am J Prev Med* 2017; 53(1):33–41. <https://doi.org/10.1016/j.amepre.2016.12.009> PMID: 28169018
34. Accordini S, Calciano L, Johannessen A, Portas L, Benediktsdóttir B, Bertelsen RJ, et al. A three-generation study of tobacco smoking with asthma. *Int J Epidemiol* 2018; 47(4):1106–1117. <https://doi.org/10.1093/ije/dyy031> PMID: 29534228
35. Breslau N, Peterson EL. Smoking cessation in young adults: age at initiation of cigarette smoking and other suspected influences. *Am J Pub Health* 1996; 86(2):214–20.
36. Chen J, Millar WJ. Age at smoking initiation: implications for quitting. *Health Rep* 1998; 9(4):39–46. PMID: 9836879
37. Kocak ND, Eren A, Boga S, Aktürk ÜA, Öztürk ÜA, Arınç S, et al. Relapse rate and factors related to relapse in a 1-year follow-up of subjects participating in a smoking cessation program. *Respir Care* 2015; 60(12):1796–803. <https://doi.org/10.4187/respcare.03883> PMID: 26286738
38. Breslau N, Fenn N, Peterson EL. Early smoking initiation and nicotine dependence in a cohort of young adults. *Drug Alcohol Dependence* 1993; 33:129–37.
39. Olivieri M, Poli A, Zuccaro P, Ferrari M, Lampronti G, de Marco R, et al. Tobacco smoke exposure and serum cotinine in a random sample of adults living in Verona, Italy. *Arch Environ Health* 2002; 57:355–9. <https://doi.org/10.1080/00039890209601421> PMID: 12530604
40. Babb S, Malarcher A, Schauer G, Asman K, Jamal A. Quitting smoking among adults—United States, 2000–2015. *MMWR Morb Mortal Wkly Rep* 2017; 65(52):1457–64. <https://doi.org/10.15585/mmwr.mm6552a1> PMID: 28056007